

时间尺度联合分数阶约束力学系统 Noether 定理^{*}

宋传静¹ 吴镇宇¹ 张毅^{2†}

(1. 苏州科技大学 数学科学学院, 苏州 215009)

(2. 苏州科技大学 土木工程学院, 苏州 215011)

摘要 研究时间尺度联合 Caputo 导数下的 Lagrange 系统和 Hamilton 系统的 Noether 定理。首先, 给出时间尺度联合 Caputo 导数的定义; 其次, 研究时间尺度联合 Caputo 导数下的系统运动微分方程, 包括 Lagrange 系统和 Hamilton 系统; 第三, 分别给出这两个系统的 Noether 恒等式和守恒量; 第四, 对于所得结果及所使用的方法, 各自举例进行说明。

关键词 时间尺度, 联合 Caputo 导数, Noether 定理, Lagrange 系统, Hamilton 系统

中图分类号:O316

文献标志码:A

Noether's Theorem of Combined Fractional Constrained Mechanical Systems on Time Scales^{*}

Song Chuanjing¹ Wu Zhenyu¹ Zhang Yi^{2†}

(1. School of Mathematical Sciences, Suzhou University of Science and Technology, Suzhou 215009, China)

(2. School of Civil Engineering, Suzhou University of Science and Technology, Suzhou 215011, China)

Abstract Noether theorems for Lagrangian and Hamiltonian systems in terms of combined Caputo derivative on time scales are investigated. Firstly, the definition of the combined Caputo derivative on time scales is given; Secondly, differential equations of motion in terms of the combined Caputo derivative on time scales are presented; Thirdly, Noether identities and conserved quantities for those two systems are studied separately; Fourthly, examples are provided.

Key words time scale, combined Caputo derivative, Noether theorem, Lagrangian system, Hamiltonian system

引言

1918 年 Emmy Noether 发表论文《不变变分问题》, 证明了 Noether 定理^[1]。Noether 定理不仅在理论物理和数学分析中具有重要地位, 还在量子

力学和广义相对论中有广泛应用, 极大地推动了科学的发展。Noether 定理的研究在 Lagrange 系统、Hamilton 系统和 Birkhoff 系统中取得了一系列成果^[2-9]。

时间尺度的概念于 1988 年被首次提出^[10-12]。

2024-11-23 收到第 1 稿, 2025-02-05 收到修改稿。

* 国家自然科学基金资助项目(12172241, 12272248), 江苏省高校“青蓝工程”资助项目, 江苏省研究生科研创新计划资助项目(KYCX24_3410), National Natural Science Foundation of China (12172241, 12272248), Qing Lan Project of Colleges and Universities in Jiangsu Province, Innovation Program for Postgraduate in Higher Education Institutions of Jiangsu Province (KYCX24_3410).

† 通信作者 E-mail: zhy@mail.usts.edu.cn

实数集、整数集、自然数集、闭区间的并集及康托尔集等均是时间尺度的例子。由于实数集和整数集可以同时被概括在时间尺度里,因此说时间尺度具有统一的特征。另一方面,时间尺度上的研究成果,可以根据需要,选择特殊的定义域进行具体化,定义域不同,所得结果也有各自的特点,因此说时间尺度具有拓展的特征。时间尺度 Noether 定理的研究始于 Bartosiewicz^[13]。时间尺度上 Noether 定理的研究涉及 Lagrange 系统^[14-16]、Hamilton 系统^[17-19]、Birkhoff 系统^[20-22]及非完整系统^[23-24]等。

分数阶微积分的提出早在 1695 年,它与整数阶微积分的提出在同一个时代。分数阶导数的类型多种多样。2015 年,Luo 和 Xu^[25]整理了比较广泛的联合分数阶导数。分数阶 Noether 定理的研究始于 Frederico 和 Torres^[26],且取得一系列研究成果^[27-33]。

时间尺度分数阶方面的研究也取得一些成果,如时间尺度分数阶原理和不等式^[34]、时间尺度分数阶动力方程^[35]、时间尺度分数阶混沌系统^[36]、时间尺度分数阶递归神经网络^[37]等。时间尺度分数阶约束力学系统对称性的研究刚刚起步,目前仅研究了时间尺度 Caputo 分数阶约束力学系统的 Noether 定理^[38-40]。考虑到联合分数阶导数的普遍性和广泛性,本文拟研究时间尺度联合分数阶约束力学系统的 Noether 定理。

文章结构安排如下:第一节列出时间尺度分数阶微积分必要的预备知识;第二节根据变分原理建立时间尺度联合 Caputo 分数阶 Lagrange 方程和时间尺度联合 Caputo 分数阶 Hamilton 方程;第三节研究 Noether 恒等式,并进一步给出时间尺度联合 Caputo 分数阶 Lagrange 系统和时间尺度联合 Caputo 分数阶 Hamilton 系统的 Noether 定理;第四节给出例题,通过举例说明本文所得结果及所使用的方法。

1 预备知识

定义 1^[41] 设 \mathbb{T} 为时间尺度,函数 $f: [a, b] \rightarrow \mathbb{R}$, $\alpha \geq 0$, 则基于时间尺度的左、右 Riemann-Liouville 积分, ${}_a^c I_{\Delta,t}^\alpha f(t)$, ${}_t^c I_{\Delta,b}^\alpha f(t)$, 分别为

$$\alpha = 0, {}_a^c I_{\Delta,t}^\alpha f(t) = f(t); \alpha > 0,$$

$${}_a^c I_{\Delta,t}^\alpha f(t) = \int_a^t h_{\alpha-1}[t, \sigma(\tau)] f(\tau) \Delta\tau \quad (1)$$

和

$$\alpha = 0, {}_t^c I_{\Delta,b}^\alpha f(t) = f(t); \alpha > 0,$$

$${}_t^c I_{\Delta,b}^\alpha f(t) = \int_t^b h_{\alpha-1}[\sigma(\tau), t] f(\tau) \Delta\tau \quad (2)$$

其中, $h_\alpha: \mathbb{T} \times \mathbb{T} \rightarrow \mathbb{R}$, $\alpha \geq 0$ 为右连续函数, 满足

$$h_{\alpha+1}(t, s) = \int_s^t h_\alpha(\tau, s) \Delta\tau, h_0(t, s) = 1, \\ t, s \in \mathbb{T} \quad (3)$$

下文将 Riemann-Liouville 简写为 R-L。

注 1 当 $\mathbb{T} = \mathbb{R}$ 时, 有 $h_{\alpha-1}(t, a) = (t-a)^{\alpha-1}/\Gamma(\alpha)$, $h_{\alpha-1}(b, t) = (b-t)^{\alpha-1}/\Gamma(\alpha)$ 时, 此时有

$${}_a^c I_t^\alpha f(t) = \int_a^t \frac{(t-\tau)^{\alpha-1}}{\Gamma(\alpha)} f(\tau) d\tau, t > a; \\ {}_t^c I_b^\alpha f(t) = \int_t^b \frac{(\tau-t)^{\alpha-1}}{\Gamma(\alpha)} f(\tau) d\tau, t < b \quad (4)$$

式(4)即为传统的左、右 R-L 积分。

定义 2^[41] 设 \mathbb{T} 是时间尺度, 函数 $f: \mathbb{T} \rightarrow \mathbb{R}$, $m-1 \leq \alpha < m$, $m \in \mathbb{N}_+$, 时间尺度左、右 R-L 导数和时间尺度左、右 Caputo 导数分别为

$${}_{RL}^a D_{\Delta,t}^\alpha f(t) = D_\Delta^m [{}_a^c I_{\Delta,t}^{m-\alpha} f(t)] \\ = D_\Delta^m \left\{ \int_a^t h_{m-\alpha-1}[t, \sigma(\tau)] f(\tau) \Delta\tau \right\} \quad (5)$$

$${}_{RL}^a D_{\Delta,b}^\alpha f(t) = (-1)^m D_\Delta^m [{}_t^c I_{\Delta,b}^{m-\alpha} f(t)] \\ = (-1)^m D_\Delta^m \left\{ \int_t^b h_{m-\alpha-1}[\sigma(\tau), t] f(\tau) \Delta\tau \right\} \quad (6)$$

$${}^C D_{\Delta,t}^\alpha f(t) = {}_a^c I_{\Delta,t}^{m-\alpha} [D_\Delta^m f(t)] = \\ \int_a^t h_{m-\alpha-1}[t, \sigma(\tau)] [D_\Delta^m f(\tau)] \Delta\tau \quad (7)$$

$${}^C D_{\Delta,b}^\alpha f(t) = (-1)^m {}_t^c I_{\Delta,b}^{m-\alpha} [D_\Delta^m f(t)] \\ = (-1)^m \int_t^b h_{m-\alpha-1}[\sigma(\tau), t] [D_\Delta^m f(\tau)] \Delta\tau \quad (8)$$

其中 D_Δ^m 表示时间尺度上求 m 阶导数。

注 2 当 $\mathbb{T} = \mathbb{R}$ 时, 式(5)~(8) 分别退化为传统的左、右 R-L 导数和左、右 Caputo 导数。

时间尺度联合 R-L 导数和时间尺度联合 Caputo 导数可定义为

$${}_{RL}^{\alpha,\beta} D_{\Delta,t}^\gamma f(t) = \gamma {}_{RL}^{\alpha,\beta} D_{\Delta,t}^\alpha f(t) + (-1)^n (1-\gamma) {}_{RL}^{\alpha,\beta} D_{\Delta,b}^\beta f(t) \quad (9)$$

$${}^C D_{\Delta,\gamma}^{\alpha,\beta} f(t) = \gamma {}^C D_{\Delta,t}^\alpha f(t) + (-1)^n (1-\gamma) {}^C D_{\Delta,b}^\beta f(t) \quad (10)$$

其中, $n-1 \leq \alpha, \beta < n, \gamma \in [0, 1]$ 。

特别地, 本文中均令 $n=1$, 则有 $0 \leq \alpha, \beta < 1$ 。

注 3 当 $\gamma=1$ 时, 由式(9)和式(10)可得

$${}_{RL}^{\alpha,\beta} D_{\Delta,t}^\gamma f(t) = {}_{RL}^{\alpha,\beta} D_{\Delta,t}^\alpha f(t), {}^C D_{\Delta,\gamma}^{\alpha,\beta} f(t) = {}^C D_{\Delta,t}^\alpha f(t) \quad (11)$$

即时间尺度联合 R-L 导数和时间尺度联合 Caputo 导数分别退化为时间尺度左 R-L 和时间尺度左 Caputo 导数. 当 $\gamma=0$ 时, 由式(9)和式(10)可得

$$\begin{aligned} {}^{\text{RL}}D_{\Delta,\gamma}^{\alpha,\beta}f(t) &= -{}^{\text{RL}}D_{\Delta,b}^{\beta}f(t), \\ {}^cD_{\Delta,\gamma}^{\alpha,\beta}f(t) &= -{}^cD_{\Delta,b}^{\beta}f(t) \end{aligned} \quad (12)$$

即时间尺度联合 R-L 导数和时间尺度联合 Caputo 导数分别退化为时间尺度右 R-L 和右 Caputo 导数. 当 $\alpha=\beta, \gamma=1/2$ 时, 由式(9)和式(10)可得

$$\begin{aligned} {}^{\text{RL}}D_{\Delta,1/2}^{\alpha,\beta}f(t) &= \frac{1}{2}{}_a^{\text{RL}}D_{\Delta,t}^{\alpha}f(t) - \frac{1}{2}{}_t^{\text{RL}}D_{\Delta,b}^{\beta}f(t) \\ &= {}_a^{\text{RL}}D_{\Delta,b}^{\alpha}f(t) \end{aligned} \quad (13)$$

$$\begin{aligned} {}^cD_{\Delta,1/2}^{\alpha,\beta}f(t) &= \frac{1}{2}{}_a^cD_{\Delta,t}^{\alpha}f(t) - \frac{1}{2}{}_t^cD_{\Delta,b}^{\beta}f(t) \\ &= {}_a^cD_{\Delta,b}^{\alpha}f(t) \end{aligned} \quad (14)$$

即时间尺度联合 R-L 导数和时间尺度联合 Caputo 导数分别退化为时间尺度 Riesz-R-L 和 Riesz-Caputo 导数. 当 $T=\mathbb{R}$ 时, 式(9)和式(10)分别退化为传统的联合 R-L 导数和联合 Caputo 导数

$$\begin{aligned} {}^{\text{RL}}D_{\gamma}^{\alpha,\beta}f(t) &= \gamma{}_a^{\text{RL}}D_t^{\alpha}f(t) + (-1)^n(1-\gamma){}_t^{\text{RL}}D_b^{\beta}f(t) \end{aligned} \quad (15)$$

$$\begin{aligned} {}^cD_{\gamma}^{\alpha,\beta}f(t) &= \gamma{}_a^cD_t^{\alpha}f(t) + (-1)^n(1-\gamma){}_t^cD_b^{\beta}f(t) \end{aligned} \quad (16)$$

式(15)和式(16)可见文献[25].

时间尺度分数阶导数的分部积分公式为^[42]

$$\begin{aligned} \int_a^b g(t) {}_a^cD_{\Delta,t}^{\alpha}f(t) \Delta t &= \int_a^b f^\sigma(t) \cdot {}_t^{\text{RL}}D_{\Delta,b}^{\alpha}g(t) \Delta t + \\ &\quad [f(t) \cdot {}_t I_{\Delta,b}^{1-\alpha}g(t)] \Big|_a^b \end{aligned} \quad (17)$$

和

$$\begin{aligned} \int_a^b g(t) {}_t^cD_{\Delta,b}^{\alpha}f(t) \Delta t &= \int_a^b f^\sigma(t) \cdot {}_a^{\text{RL}}D_{\Delta,t}^{\alpha}g(t) \Delta t - \\ &\quad [f(t) \cdot {}_a I_{\Delta,t}^{1-\alpha}g(t)] \Big|_a^b \end{aligned} \quad (18)$$

2 时间尺度联合 Caputo 分数阶运动微分方程

2.1 Lagrange 系统运动微分方程

时间尺度分数阶 Lagrange 函数为 $L(t, q_k^\sigma, {}^cD_{\Delta,\gamma}^{\alpha,\beta}q_k)$, 则 Hamilton 作用量可表示为

$$S_L[q(\cdot)] = \int_a^b L(t, q_k^\sigma, {}^cD_{\Delta,\gamma}^{\alpha,\beta}q_k) \Delta t \quad (19)$$

其中 $q_k^\sigma(t) = (q_k \circ \sigma)(t), t \in [a, b], 0 < \alpha, \beta < 1, \gamma \in [0, 1]$.

时间尺度联合 Caputo 分数阶变分原理为

$$\delta S_L = 0, \delta {}_a^cD_{\Delta,t}^{\alpha}q_k = {}_a^cD_{\Delta,t}^{\alpha}\delta q_k,$$

$$\begin{aligned} \delta {}_t^{\text{RL}}D_{\Delta,b}^{\beta}q_k &= {}_t^{\text{RL}}D_{\Delta,b}^{\beta}\delta q_k, (\delta q_k)^\sigma = \delta q_k^\sigma, \\ \delta q_k(t)|_{t=a} &= \delta q_k(t)|_{t=b} = 0 \end{aligned} \quad (20)$$

其中 δ 表示等时变分.

把式(19)代入式(20)可得

$$\delta S_L = \int_a^b (\partial_j L \cdot \delta q_j^\sigma + \partial_{n+j} L \cdot \delta {}_a^cD_{\Delta,t}^{\alpha}q_j) \Delta t = 0 \quad (21)$$

其中 $\partial_j L$ 表示对 Lagrange 函数 L 的第 j 个元素求偏导数, $\partial_{n+j} L$ 表示对 Lagrange 函数 L 的第 $n+j$ 个元素求偏导数, $j = 1, 2, \dots, n$.

利用分部积分公式(17)和公式(18)可得

$$\begin{aligned} &\int_a^b \partial_{n+j} L \cdot \delta {}_a^cD_{\Delta,t}^{\alpha}q_j \Delta t \\ &= \int_a^b \partial_{n+j} L \cdot \delta [\gamma {}_a^cD_{\Delta,t}^{\alpha}q_j + (-1)^n(1-\gamma){}_t^cD_{\Delta,b}^{\beta}q_j] \Delta t \\ &= - \int_a^b \delta q_j^\sigma [(1-\gamma){}_a^cD_{\Delta,t}^{\alpha} \partial_{n+j} L - \gamma {}_t^cD_{\Delta,b}^{\beta} \partial_{n+j} L] \Delta t \\ &= - \int_a^b (\delta q_j^\sigma \cdot {}_a^{\text{RL}}D_{\Delta,1-\gamma}^{\beta,\alpha} \partial_{n+j} L) \Delta t \end{aligned} \quad (22)$$

将式(22)代入式(21)且利用式(20)中的边界条件可得

$$\begin{aligned} 0 = \delta S_L &= \int_a^b (\partial_j L \cdot \delta q_j^\sigma - \delta q_j^\sigma \cdot {}_a^{\text{RL}}D_{\Delta,1-\gamma}^{\beta,\alpha} \partial_{n+j} L) \Delta t \\ &= \int_a^b \left\{ \left[\int_a^t (\partial_j L - {}_a^{\text{RL}}D_{\Delta,1-\gamma}^{\beta,\alpha} \partial_{n+j} L) \Delta \theta \cdot \delta q_j \right]^\Delta - \right. \\ &\quad \left. (\delta q_j)^\Delta \cdot \left[\int_a^t (\partial_j L - {}_a^{\text{RL}}D_{\Delta,1-\gamma}^{\beta,\alpha} \partial_{n+j} L) \Delta \theta \right] \Delta \theta \cdot (\delta q_j)^\Delta \right\} \Delta t \\ &= - \int_a^b \left[\int_a^t (\partial_j L - {}_a^{\text{RL}}D_{\Delta,1-\gamma}^{\beta,\alpha} \partial_{n+j} L) \Delta \theta \cdot (\delta q_j)^\Delta \right] \Delta t \end{aligned} \quad (23)$$

从而

$$\int_a^t (\partial_j L - {}_a^{\text{RL}}D_{\Delta,1-\gamma}^{\beta,\alpha} \partial_{n+j} L) \Delta \theta = \text{const.} \quad (24)$$

式(24)两边同时对 t 求 Δ 导数, 可得

$$\partial_j L - {}_a^{\text{RL}}D_{\Delta,1-\gamma}^{\beta,\alpha} \partial_{n+j} L = 0 \quad (25)$$

式(25)称为时间尺度联合 Caputo 分数阶 Lagrange 方程.

注 4 当 $T=\mathbb{R}$ 时, 式(25)退化为

$$\begin{aligned} \partial_j L(t, q_k, {}^cD_{\gamma}^{\alpha,\beta}q_k) - \\ {}_a^{\text{RL}}D_{\Delta,1-\gamma}^{\beta,\alpha} \partial_{n+j} L(t, q_k, {}^cD_{\gamma}^{\alpha,\beta}q_k) = 0 \end{aligned} \quad (26)$$

式(26)为联合 Caputo 分数阶 Lagrange 方程, 与文献[43]所得结果一致.

注 5 当 $\alpha \rightarrow 1, \beta \rightarrow 1$ 时, 式(25)退化为

$$\partial_j L(t, q_k^\sigma, q_k^\Delta) - [\partial_{n+j} L(t, q_k^\sigma, q_k^\Delta)]^\Delta = 0 \quad (27)$$

式(27)为时间尺度上的 Lagrange 方程, 与文献[13]所得结果一致.

注 6 当 $\gamma=1$ 时, 式(25)退化为

$$\begin{aligned} \partial_j L(t, q_k^\sigma, {}^C D_{\Delta,t}^\alpha q_k) + \\ {}_t^{\text{RL}} D_{\Delta,t}^\alpha \partial_{n+j} L(t, q_k^\sigma, {}^C D_{\Delta,t}^\alpha q_k) = 0 \end{aligned} \quad (28)$$

式(28)为时间尺度左 Caputo 分数阶 Lagrange 方程, 其与文献[38]中所得结果一致. 当 $\gamma=0$ 时, 式(25)退化为

$$\begin{aligned} \partial_j L(t, q_k^\sigma, {}^C D_{\Delta,b}^\alpha q_k) + \\ {}_a^R D_{\Delta,t}^\alpha \partial_{n+j} L(t, q_k^\sigma, {}^C D_{\Delta,b}^\alpha q_k) = 0 \end{aligned} \quad (29)$$

式(29)称为时间尺度右 Caputo 分数阶 Lagrange 方程. 当 $\alpha=\beta, \gamma=1/2$ 时, 式(25)退化为

$$\begin{aligned} \partial_j L(t, q_k^\sigma, {}^{RC} D_{\Delta,b}^\alpha q_k) - \\ {}_a^R D_{\Delta,b}^\alpha \partial_{n+j} L(t, q_k^\sigma, {}^{RC} D_{\Delta,b}^\alpha q_k) = 0 \end{aligned} \quad (30)$$

式(30)称为时间尺度 Riesz-Caputo 分数阶 Lagrange 方程.

2.2 Hamilton 系统运动微分方程

时间尺度 Caputo 分数阶 Lagrange 函数为 $L(t, q_k^\sigma, {}^C D_{\Delta,\gamma}^{\alpha,\beta} q_k)$, 则广义动量和 Hamilton 函数为

$$\begin{aligned} p_k &= \frac{\partial L}{\partial {}^C D_{\Delta,\gamma}^{\alpha,\beta} q_k}, \\ H &= H(t, q_k^\sigma, p_k) \\ &= p_k \cdot {}^C D_{\Delta,\gamma}^{\alpha,\beta} q_k - L(t, q_k^\sigma, {}^C D_{\Delta,\gamma}^{\alpha,\beta} q_k) \end{aligned} \quad (31)$$

Hamilton 作用量可表示为

$$S_H = \int_a^b (p_k \cdot {}^C D_{\Delta,\gamma}^{\alpha,\beta} q_k - H) \Delta t \quad (32)$$

时间尺度联合 Caputo 分数阶变分原理为

$$\begin{aligned} \delta S_H &= 0, \delta {}^C D_{\Delta,t}^\alpha q_k = {}_a^C D_{\Delta,t}^\alpha \delta q_k, \\ \delta {}_t^C D_{\Delta,b}^\alpha q_k &= {}_t^C D_{\Delta,b}^\alpha \delta q_k, (\delta q_k)^\sigma = \delta q_k^\sigma, \\ \delta q_k(t)|_{t=a} &= \delta q_k(t)|_{t=b} = 0 \end{aligned} \quad (33)$$

又

$$\begin{aligned} \int_a^b p_k \cdot {}^C D_{\Delta,\gamma}^{\alpha,\beta} q_k \Delta t &= \int_a^b p_k \cdot {}^C D_{\Delta,\gamma}^{\alpha,\beta} \delta q_k \Delta t \\ &= \int_a^b p_k \cdot [\gamma {}_a^C D_{\Delta,t}^\alpha \delta q_k - (1-\gamma) {}_t^C D_{\Delta,b}^\alpha \delta q_k] \Delta t \\ &= \gamma \left[\int_a^b \delta q_k^\sigma \cdot {}_t^{\text{RL}} D_{\Delta,b}^\alpha p_k \Delta t + (\delta q_k \cdot {}_t I_{\Delta,t}^{1-\alpha} p_k) \Big|_{t=a}^{t=b} \right] - \\ &\quad (1-\gamma) \left[\int_a^b \delta q_k^\sigma \cdot {}_a^{\text{RL}} D_{\Delta,t}^\alpha p_k \Delta t - (\delta q_k \cdot {}_a I_{\Delta,t}^{1-\alpha} p_k) \Big|_{t=a}^{t=b} \right] \\ &= - \int_a^b (\delta q_k^\sigma \cdot {}_a^{\text{RL}} D_{\Delta,t}^\alpha p_k) \Delta t \end{aligned} \quad (34)$$

故

$$\begin{aligned} 0 &= \delta S_H = \int_a^b (\delta p_k \cdot {}^C D_{\Delta,\gamma}^{\alpha,\beta} q_k + p_k \cdot \delta {}^C D_{\Delta,\gamma}^{\alpha,\beta} q_k - \\ &\quad \partial_j H \cdot \delta q_k^\sigma - \partial_{n+j} H \cdot \delta p_k) \Delta t \end{aligned}$$

$$\begin{aligned} &= \int_a^b (\delta p_k \cdot {}^C D_{\Delta,\gamma}^{\alpha,\beta} q_k - \delta q_k^\sigma \cdot {}_a^{\text{RL}} D_{\Delta,t}^\alpha p_k - \\ &\quad \partial_j H \cdot \delta q_k^\sigma - \partial_{n+j} H \cdot \delta p_k) \Delta t \\ &= \int_a^b [\delta p_k \cdot ({}^C D_{\Delta,\gamma}^{\alpha,\beta} q_k - \partial_{n+j} H) - \\ &\quad \delta q_k^\sigma ({}_a^{\text{RL}} D_{\Delta,t}^\alpha p_k + \partial_j H)] \Delta t \end{aligned} \quad (35)$$

将式(31)中的 Hamilton 函数两边对 p_j 求导, 可得

$$\partial_{n+j} H = {}^C D_{\Delta,\gamma}^{\alpha,\beta} q_j \quad (36)$$

将式(36)代入式(35)可得

$$\begin{aligned} 0 &= \delta S_H = - \int_a^b \delta q_k^\sigma ({}_a^{\text{RL}} D_{\Delta,t}^\alpha p_k + \partial_j H) \Delta t \\ &= - \int_a^b \{ [\int_a^t ({}_a^{\text{RL}} D_{\Delta,t}^\alpha p_k + \partial_j H) \Delta \theta \cdot \delta q_k]^\Delta - \\ &\quad (\delta q_k)^\Delta \int_a^t ({}_a^{\text{RL}} D_{\Delta,t}^\alpha p_k + \partial_j H) \Delta \theta \} \Delta t \\ &= \int_a^b [(\delta q_k)^\Delta \cdot \int_a^t ({}_a^{\text{RL}} D_{\Delta,t}^\alpha p_k + \partial_j H) \Delta \theta] \Delta t \end{aligned} \quad (37)$$

从而

$$\int_a^t ({}_a^{\text{RL}} D_{\Delta,t}^\alpha p_k + \partial_j H) \Delta \theta = \text{const.} \quad (38)$$

即

$${}_a^{\text{RL}} D_{\Delta,t}^\alpha p_k + \partial_j H = 0 \quad (39)$$

将式(36)与式(39)放在一起可得

$$\begin{cases} \partial_j H = - {}_a^{\text{RL}} D_{\Delta,t}^\alpha p_k \\ \partial_{n+j} H = {}^C D_{\Delta,\gamma}^{\alpha,\beta} q_j \end{cases} \quad (40)$$

式(40)称为时间尺度联合 Caputo 分数阶 Hamilton 方程.

注 7 当 $T=\mathbb{R}$ 时, $H=H(t, q_k, p_k)$ 时, 式(40)退化为

$$\partial_j H = - {}_a^{\text{RL}} D_{\Delta,t}^\alpha p_k, \partial_{n+j} H = {}^C D_{\Delta,\gamma}^{\alpha,\beta} q_j \quad (41)$$

式(41)为联合 Caputo 分数阶 Hamilton 方程, 其与文献[43]中所得结果一致.

注 8 当 $\alpha \rightarrow 1, \beta \rightarrow 1$ 时, $H=H(t, q_k^\sigma, p_k)$, 此时式(40)退化为

$$\partial_j H = - p_j^\Delta, \partial_{n+j} H = q_j^\Delta \quad (42)$$

式(42)为时间尺度上的 Hamilton 方程, 其与文献[18]中所得结果一致.

注 9 当 $\gamma=1$ 时, 方程(40)退化为

$$\begin{cases} \partial_j H(t, q_k^\sigma, p_k) = {}_a^{\text{RL}} D_{\Delta,t}^\alpha p_k \\ \partial_{n+j} H(t, q_k^\sigma, p_k) = {}^C D_{\Delta,\gamma}^{\alpha,\beta} q_j \end{cases} \quad (43)$$

式(43)为时间尺度左 Caputo 分数阶 Hamilton 方程, 其与文献[39]所得结果一致. 当 $\gamma=0$ 时, 式(40)退化为

$$\begin{aligned}\partial_j H(t, q_k^\sigma, p_k) &= -_a^{\text{RL}} D_{\Delta, t}^\alpha p_j, \\ \partial_{n+j} H(t, q_k^\sigma, p_k) &= {}_t^C D_{\Delta, b}^\alpha q_j\end{aligned}\quad (44)$$

式(44)称为时间尺度右 Caputo 分数阶 Hamilton 方程. 当 $\alpha = \beta, \gamma = 1/2$ 时, 方程(40)退化为

$$\begin{aligned}\partial_j H(t, q_k^\sigma, p_k) &= -_a^R D_{\Delta, t}^\alpha p_j, \\ \partial_{n+j} H(t, q_k^\sigma, p_k) &= {}_a^{\text{RC}} D_{\Delta, b}^\alpha q_j\end{aligned}\quad (45)$$

式(45)称为时间尺度 Riesz-Caputo 分数阶 Hamilton 方程.

3 时间尺度联合 Caputo 分数阶 Noether 定理

3.1 Lagrange 系统 Noether 定理

定义 3 对于一个物理量 I , 如果满足 $\Delta I / \Delta t = 0$, 则称 I 为一个守恒量.

引入无限小变换

$$\begin{aligned}\bar{t} &= T_{L_\varepsilon} = t + \varepsilon \xi_{L_0} + o(\varepsilon), \\ \bar{q}_j &= Q_{L_\varepsilon}^j = q_j + \varepsilon \xi_{L_j} + o(\varepsilon), \quad j = 1, 2, \dots, n\end{aligned}\quad (46)$$

其中 ξ_{L_0}, ξ_{L_j} 是无限小生成元, ε 是无限小参数. 设映射 $t \mapsto \nu(t); T_{L_\varepsilon}(t, q_k) \in \mathbb{R}$ 为递增的, 映射的像构成一个新的时间尺度 \mathbb{T}' , 其前跳算子为 $\bar{\sigma}$, 对应导数为 $\bar{\Delta}$, 且满足 $\sigma \circ \nu = \nu \circ \sigma$.

Hamilton 作用量的不变性可表示为, 对于任意的 $[t_a, t_b] \subset [a, b]$, 恒有

$$\begin{aligned}&\int_{t_a}^{t_b} L[t, q_k^\sigma(t), {}^C D_{\Delta, \gamma}^\alpha q_k(t)] \Delta t \\ &= \int_{t_a}^{t_b} L[\bar{t}, \bar{q}_k^\sigma(\bar{t}), {}^C D_{\Delta, \gamma}^\alpha \bar{q}_k(\bar{t})] \bar{\Delta} t\end{aligned}\quad (47)$$

成立. 即

$$\begin{aligned}&\int_{t_a}^{t_b} L[t, q_k^\sigma(t), {}^C D_{\Delta, \gamma}^\alpha q_k(t)] \Delta t \\ &= \int_{t_a}^{t_b} L \left[T_{L_\varepsilon}, (Q_{L_\varepsilon}^k)^\sigma, \frac{{}^C D_{\Delta, \gamma}^\alpha Q_{L_\varepsilon}^k}{(T_{L_\varepsilon}^\Delta)^a} \right] \cdot T_{L_\varepsilon}^\Delta \Delta t\end{aligned}\quad (48)$$

其中

$${}^C D_{\Delta, \gamma}^\alpha \bar{q}_k(\bar{t}) = \frac{{}^C D_{\Delta, \gamma}^\alpha (\bar{q}_k \circ \nu)(t)}{[v^\Delta(t)]^a}\quad (49)$$

式(48)等价于

$$\begin{aligned}&L[t, q_k^\sigma(t), {}^C D_{\Delta, \gamma}^\alpha q_k(t)] \\ &= L \left[T_{L_\varepsilon}, (Q_{L_\varepsilon}^k)^\sigma, \frac{{}^C D_{\Delta, \gamma}^\alpha Q_{L_\varepsilon}^k}{(T_{L_\varepsilon}^\Delta)^a} \right] \cdot T_{L_\varepsilon}^\Delta\end{aligned}\quad (50)$$

即

$$0 = \partial_0 L \cdot \xi_{L_0} + \partial_k L \cdot \xi_{L_k}^\sigma + \partial_{n+k} L \cdot$$

$$\begin{aligned}\frac{{}^C D_{\Delta, \gamma}^\alpha \xi_{L_k} - \alpha {}^C D_{\Delta, \gamma}^\alpha q_k \cdot \xi_{L_0}^\Delta}{(T_{L_\varepsilon}^\Delta)^{2a}} \Big|_{\varepsilon=0} &+ L \xi_{L_0}^\Delta \\ &= \partial_0 L \cdot \xi_{L_0} + \partial_k L \cdot \xi_{L_k}^\sigma + \partial_{n+k} L \cdot ({}^C D_{\Delta, \gamma}^\alpha \xi_{L_k} - \\ &\quad \alpha {}^C D_{\Delta, \gamma}^\alpha q_k \cdot \xi_{L_0}^\Delta) + L \xi_{L_0}^\Delta\end{aligned}\quad (51)$$

从而 Noether 恒等式为

$$\begin{aligned}\partial_0 L \cdot \xi_{L_0} + \partial_k L \cdot \xi_{L_k}^\sigma + \partial_{n+k} L \cdot ({}^C D_{\Delta, \gamma}^\alpha \xi_{L_k} - \\ \alpha {}^C D_{\Delta, \gamma}^\alpha q_k \cdot \xi_{L_0}^\Delta) + L \xi_{L_0}^\Delta &= 0\end{aligned}\quad (52)$$

定理 1 对于时间尺度联合 Caputo 分数阶 Lagrange 系统[式(25)], 如果 ξ_{L_0} 和 ξ_{L_j} 满足 Noether 恒等式[式(52)], 则该系统存在如下守恒量

$$\begin{aligned}I_L = \int_a^t [\partial_0 L + (\alpha \cdot \partial_{n+k} L \cdot {}^C D_{\Delta, \gamma}^\alpha q_k + \partial_0 L \cdot \mu(\tau) - \\ L)^\Delta] \cdot \xi_{L_0}^\sigma \Delta \tau + [L - \alpha \cdot \partial_{n+k} L \cdot {}^C D_{\Delta, \gamma}^\alpha q_k - \partial_0 L \cdot \\ \mu(\tau)] \cdot \xi_{L_0} + \int_a^t ({}^{\text{RL}} D_{\Delta, 1-\gamma}^{\beta, \alpha} \partial_{n+k} L \cdot \xi_{L_k}^\sigma + \\ \partial_{n+k} L \cdot {}^C D_{\Delta, \gamma}^\alpha \xi_{L_k}) \Delta \tau\end{aligned}\quad (53)$$

证明 利用式(25)和式(52)可得

$$\begin{aligned}\frac{\Delta}{\Delta t} I_L &= \partial_0 L \cdot \xi_{L_0} + [\alpha \cdot \partial_{n+k} L \cdot {}^C D_{\Delta, \gamma}^\alpha q_k + \partial_0 L \cdot \\ \mu(t) - L]^\Delta \cdot \xi_{L_0} + [L - \alpha \cdot \partial_{n+k} L \cdot {}^C D_{\Delta, \gamma}^\alpha q_k - \\ \partial_0 L \cdot \mu(t)] \cdot \xi_{L_0} + [L - \alpha \cdot \partial_{n+k} L \cdot {}^C D_{\Delta, \gamma}^\alpha q_k - \\ \partial_0 L \cdot \mu(t)]^\Delta \cdot \xi_{L_0} + {}^{\text{RL}} D_{\Delta, 1-\gamma}^{\beta, \alpha} \partial_{n+k} L \cdot \xi_{L_k} + \partial_{n+k} L \cdot \\ {}^C D_{\Delta, \gamma}^\alpha \xi_{L_k} = -\partial_k L \cdot \xi_{L_k}^\sigma + {}^{\text{RL}} D_{\Delta, 1-\gamma}^{\beta, \alpha} \partial_{n+k} L \cdot \xi_{L_k} = 0\end{aligned}$$

证毕.

注 10 当 $\mathbb{T} = \mathbb{R}$ 时, 式(52)退化为

$$\begin{aligned}\partial_0 L \cdot \xi_{L_0} + \partial_k L \cdot \xi_{L_k}^\sigma + \partial_{n+k} L \cdot ({}^C D_{\gamma}^{\alpha, \beta} \xi_{L_k} - \\ \alpha {}^C D_{\gamma}^{\alpha, \beta} q_k \cdot \dot{\xi}_{L_0}) + L \dot{\xi}_{L_0} = 0\end{aligned}\quad (54)$$

式(53)退化为

$$\begin{aligned}I_L = \int_a^t \left[\partial_0 L + \frac{d}{dt} (\alpha \cdot \partial_{n+k} L \cdot {}^C D_{\gamma}^{\alpha, \beta} q_k - L) \right] \cdot \\ \xi_{L_0} d\tau + (L - \alpha \cdot \partial_{n+k} L \cdot {}^C D_{\gamma}^{\alpha, \beta} q_k) \cdot \xi_{L_0} + \\ \int_a^t ({}^{\text{RL}} D_{1-\gamma}^{\beta, \alpha} \partial_{n+k} L \cdot \xi_{L_k} + \partial_{n+k} L \cdot {}^C D_{\gamma}^{\alpha, \beta} \xi_{L_k}) d\tau\end{aligned}\quad (55)$$

定理 2 对于联合 Caputo 分数阶 Lagrange 系统[式(26)], 如果 ξ_{L_0} 和 ξ_{L_j} 满足式(54), 则该系统存在如式(55)的守恒量.

证明 利用式(26)和式(54)可得

$$\begin{aligned}\frac{d}{dt} I_L &= \partial_0 L \cdot \xi_{L_0} + \frac{d}{dt} (\alpha \cdot \partial_{n+k} L \cdot {}^C D_{\gamma}^{\alpha, \beta} q_k - L) \cdot \\ \xi_{L_0} + (L - \alpha \cdot \partial_{n+k} L \cdot {}^C D_{\gamma}^{\alpha, \beta} q_k) \dot{\xi}_{L_0} + \frac{d}{dt} (L - \\ \alpha \cdot \partial_{n+k} L \cdot {}^C D_{\gamma}^{\alpha, \beta} q_k) \cdot \xi_{L_0} + {}^{\text{RL}} D_{1-\gamma}^{\beta, \alpha} \partial_{n+k} L \cdot \xi_{L_k} + \\ \partial_{n+k} L \cdot {}^C D_{\gamma}^{\alpha, \beta} \xi_{L_k} = (-\partial_k L + {}^{\text{RL}} D_{1-\gamma}^{\beta, \alpha} \partial_{n+k} L) \cdot\end{aligned}$$

$$\xi_{Lk} = 0$$

证毕.

注 11 当 $\alpha \rightarrow 1, \beta \rightarrow 1$ 时, 式(52)退化为

$$\begin{aligned} & \partial_0 L \cdot \xi_{L0} + \partial_k L \cdot \xi_{Lk}^\sigma + \partial_{k+n} L \cdot \\ & (\xi_{Lk}^\Delta - q_k^\Delta \cdot \xi_{L0}^\Delta) + L \xi_{L0}^\Delta = 0 \end{aligned} \quad (56)$$

式(53)退化为

$$\begin{aligned} I_L = & \int_a^t \{ \partial_0 L + [\partial_{n+k} L \cdot q_k^\Delta + \partial_0 L \cdot \mu(\tau) - L]^\Delta \cdot \\ & \xi_{L0}^\sigma \Delta\tau + [L - \partial_{n+k} L \cdot q_k^\Delta - \partial_0 L \cdot \mu(t)] \cdot \\ & \xi_{L0} + \partial_{n+k} L \cdot \xi_{Lk} \} \Delta\tau \end{aligned} \quad (57)$$

定理 3 对于时间尺度上的 Lagrange 系统[式(27)], 如果 ξ_{L0} 和 ξ_{Lj} 满足式(56), 则该系统存在如式(57)的守恒量.

证明 利用式(27)和式(56)可得 $\Delta I_L / \Delta t = 0$.

证毕. 此定理与文献[44]所得结果一致.

注 12 当 $\gamma=1$ 时, 可得时间尺度左 Caputo 分数阶 Lagrange 系统[式(28)], 其 Noether 恒等式为

$$\begin{aligned} & \partial_0 L \cdot \xi_{L0} + \partial_k L \cdot \xi_{Lk}^\sigma + \partial_{k+n} L \cdot ({}^C D_{\Delta,t}^\alpha \xi_{Lk} - \\ & \alpha {}^C D_{\Delta,t}^\alpha q_k \cdot \xi_{L0}^\Delta) + L \xi_{L0}^\Delta = 0 \end{aligned} \quad (58)$$

对应 Noether 守恒量为

$$\begin{aligned} I_L = & \int_a^t \{ \partial_0 L + [\alpha \cdot \partial_{n+k} L \cdot {}^C D_{\Delta,t}^\alpha q_k + \partial_0 L \cdot \mu(\tau) - \\ & L]^\Delta \} \cdot \xi_{L0}^\sigma \Delta\tau + [L - \alpha \cdot \partial_{n+k} L \cdot {}^C D_{\Delta,t}^\alpha q_k - \partial_0 L \cdot \\ & \mu(t)] \cdot \xi_{L0} + \int_a^t (-{}^R D_{\Delta,b}^\alpha \partial_{n+k} L \cdot \xi_{Lk}^\sigma + \\ & \partial_{n+k} L \cdot {}^C D_{\Delta,t}^\alpha \xi_{Lk}) \Delta\tau \end{aligned} \quad (59)$$

此结果与文献[38]所得结果一致.

注 13 当 $\gamma=0$ 时, 可得时间尺度右 Caputo 分数阶 Lagrange 系统[式(29)], 其 Noether 恒等式为

$$\begin{aligned} & \partial_0 L \cdot \xi_{L0} + \partial_k L \cdot \xi_{Lk}^\sigma + \partial_{k+n} L \cdot (-{}^C D_{\Delta,b}^\alpha \xi_{Lk} + \\ & \alpha {}^C D_{\Delta,b}^\alpha q_k \cdot \xi_{L0}^\Delta) + L \xi_{L0}^\Delta = 0 \end{aligned} \quad (60)$$

对应 Noether 守恒量为

$$\begin{aligned} I_L = & \int_a^t \{ \partial_0 L + [-\alpha \cdot \partial_{n+k} L \cdot {}^C D_{\Delta,b}^\alpha q_k + \partial_0 L \cdot \\ & \mu(\tau) - L]^\Delta \} \cdot \xi_{L0}^\sigma \Delta\tau + [L + \alpha \cdot \partial_{n+k} L \cdot \\ & {}^C D_{\Delta,b}^\alpha q_k - \partial_0 L \cdot \mu(t)] \cdot \xi_{L0} + \int_a^t ({}^R D_{\Delta,t}^\alpha \partial_{n+k} L \cdot \\ & \xi_{Lk}^\sigma - \partial_{n+k} L \cdot {}^C D_{\Delta,b}^\alpha \xi_{Lk}) \Delta\tau \end{aligned} \quad (61)$$

注 14 当 $\alpha=\beta, \gamma=1/2$ 时, 可得时间尺度 Riesz-Caputo 分数阶 Lagrange 系统[式(30)], 其 Noether 恒等式为

$$\begin{aligned} & \partial_0 L \cdot \xi_{L0} + \partial_k L \cdot \xi_{Lk}^\sigma + \partial_{k+n} L \cdot ({}^C D_{\Delta,b}^\alpha \xi_{Lk} - \\ & \alpha {}^C D_{\Delta,b}^\alpha q_k \cdot \xi_{L0}^\Delta) + L \xi_{L0}^\Delta = 0 \end{aligned} \quad (62)$$

对应 Noether 守恒量为

$$\begin{aligned} I_L = & \int_a^t \{ \partial_0 L + [\alpha \cdot \partial_{n+k} L \cdot {}^C D_{\Delta,b}^\alpha q_k + \partial_0 L \cdot \\ & \mu(\tau) - L]^\Delta \} \cdot \xi_{L0}^\sigma \Delta\tau + [L - \alpha \cdot \partial_{n+k} L \cdot \\ & {}^C D_{\Delta,b}^\alpha q_k - \partial_0 L \cdot \mu(t)] \cdot \xi_{L0} + \int_a^t ({}^R D_{\Delta,t}^\alpha \partial_{n+k} L \cdot \\ & \xi_{Lk}^\sigma + \partial_{n+k} L \cdot {}^C D_{\Delta,b}^\alpha \xi_{Lk}) \Delta\tau \end{aligned} \quad (63)$$

3.2 Hamilton 系统 Noether 定理

取无限小变换为

$$\begin{aligned} \bar{t} &= T_{H\epsilon} = t + \epsilon \xi_{H0} + o(\epsilon), \\ \bar{q}_j &= Q_{H\epsilon}^j = q_j + \epsilon \xi_{Hj} + o(\epsilon), \\ \bar{p}_j &= P_{H\epsilon}^j = p_j + \epsilon \eta_{Hj} + o(\epsilon) \end{aligned} \quad (64)$$

其中 $\xi_{H0}, \xi_{Hj}, \eta_{Hj}, j=1, 2, \dots, n$ 是无限小生成元, ϵ 是无限小参数. 设映射 $t \mapsto \nu(t): T_{H\epsilon}(t, q_k, p_k) \in \mathbb{R}$ 为递增的, 映射的像构成一个新的时间尺度 \mathbb{T}' , 其前跳算子为 $\bar{\sigma}$, 对应导数为 $\bar{\Delta}$, 且满足 $\sigma \circ \nu = \nu \circ \sigma$.

Hamilton 作用量的不变性可表示为对于任意的 $[t_a, t_b] \subset [a, b]$, 恒有

$$\begin{aligned} & \int_{t_a}^{t_b} \{ p_j(t) \cdot {}^C D_{\Delta,\gamma}^{\alpha,\beta} q_j(t) - H[t, q_k^\sigma(t), p_k(t)] \} \Delta t \\ & = \int_{\bar{t}_a}^{\bar{t}_b} \{ \bar{p}_j(\bar{t}) \cdot {}^C D_{\Delta,\gamma}^{\alpha,\beta} \bar{q}_j(\bar{t}) - \bar{H}[\bar{t}, \bar{q}_k^\sigma(\bar{t}), \bar{p}_k(\bar{t})] \} \Delta \bar{t} \end{aligned} \quad (65)$$

成立. 利用式(49)可得

$$\begin{aligned} & \int_{t_a}^{t_b} \{ p_j(t) \cdot {}^C D_{\Delta,\gamma}^{\alpha,\beta} q_j(t) - H[t, q_k^\sigma(t), p_k(t)] \\ & p_k(t) \} \Delta t \\ & = \int_{t_a}^{t_b} \left\{ P_{H\epsilon}^j \cdot \frac{{}^C D_{\Delta,\gamma}^{\alpha,\beta} Q_{H\epsilon}^j}{(T_{H\epsilon}^\Delta)^a} - H[T_{H\epsilon}, (Q_{H\epsilon}^k)^\sigma, P_{H\epsilon}^k] \right\} T_{H\epsilon}^\Delta \Delta t \end{aligned} \quad (66)$$

由 $[t_a, t_b]$ 的任意性可得

$$\begin{aligned} & p_j(t) \cdot {}^C D_{\Delta,\gamma}^{\alpha,\beta} q_j(t) - H[t, q_k^\sigma(t), p_k(t)] \\ & = \left\{ P_{H\epsilon}^j \cdot \frac{{}^C D_{\Delta,\gamma}^{\alpha,\beta} Q_{H\epsilon}^j}{(T_{H\epsilon}^\Delta)^a} - H[T_{H\epsilon}, (Q_{H\epsilon}^k)^\sigma, P_{H\epsilon}^k] \right\} T_{H\epsilon}^\Delta \end{aligned} \quad (67)$$

即

$$\begin{aligned} & p_j \cdot {}^C D_{\Delta,\gamma}^{\alpha,\beta} \xi_{Hj} - (\alpha - 1) p_j \cdot \xi_{H0}^\Delta {}^C D_{\Delta,\gamma}^{\alpha,\beta} q_j - \\ & H \xi_{H0}^\Delta - \partial_0 H \cdot \xi_{H0} - \partial_k H \cdot \xi_{Hk}^\sigma = 0 \end{aligned} \quad (68)$$

式(68)称为时间尺度联合 Caputo 分数阶 Noether 恒等式.

定理 4 对于时间尺度联合 Caputo 分数阶 Hamilton 系统[式(40)], 若无限小生成元满足式(68), 则该系统存在如下形式的守恒量

$$\begin{aligned} I_H = & \int_a^t \{ [H + (\alpha - 1)p_k \cdot \partial_{n+k} H - \partial_0 H \cdot \mu(\tau)]^\Delta - \\ & \partial_0 H \} \cdot \xi_{H0}^\sigma \Delta\tau - [H + (\alpha - 1)p_k \cdot \partial_{n+k} H - \\ & \partial_0 H \cdot \mu(t)] \cdot \xi_{H0} + \int_a^t (p_k \cdot {}^C D_{\Delta,\gamma}^{\alpha,\beta} \xi_{Hk} + \\ & \xi_{Hk}^\sigma \cdot {}^{RL} D_{\Delta,1-\gamma}^{\beta,\alpha} p_k) \Delta\tau \end{aligned} \quad (69)$$

证明 利用式(40)和式(68)可得

$$\begin{aligned} \frac{\Delta I_H}{\Delta t} = & [H + (\alpha - 1)p_k \cdot \partial_{n+k} H - \partial_0 H \cdot \mu(\tau)]^\Delta \xi_{H0} - \\ & \partial_0 H \cdot \xi_{H0}^\sigma - [H + (\alpha - 1)p_k \cdot \partial_{n+k} H - \partial_0 H \cdot \\ & \mu(t)]^\Delta \cdot \xi_{H0}^\sigma - [H + (\alpha - 1)p_k \cdot \partial_{n+k} H - \partial_0 H \cdot \\ & \mu(t)] \cdot \xi_{H0}^\Delta + p_k \cdot {}^C D_{\Delta,\gamma}^{\alpha,\beta} \xi_{Hk} + \xi_{Hk}^\sigma \cdot {}^{RL} D_{\Delta,1-\gamma}^{\beta,\alpha} p_k \\ = & \partial_k H \cdot \xi_{Hk}^\sigma + \xi_{Hk}^\sigma \cdot {}^{RL} D_{\Delta,1-\gamma}^{\beta,\alpha} p_k \\ = & \xi_{Hk}^\sigma (\partial_k H + {}^{RL} D_{\Delta,1-\gamma}^{\beta,\alpha} p_k) = 0 \end{aligned}$$

证毕.

注 15 当 $\mathbb{T} = \mathbb{R}$ 时, 式(68)退化为

$$\begin{aligned} p_j \cdot {}^C D_\gamma^{\alpha,\beta} \xi_{Hj} - (\alpha - 1)p_j \cdot \dot{\xi}_{H0} {}^C D_\gamma^{\alpha,\beta} q_j - \\ H \dot{\xi}_{H0} - \partial_0 H \cdot \xi_{H0} - \partial_k H \cdot \xi_{Hk} = 0 \end{aligned} \quad (70)$$

式(69)退化为

$$\begin{aligned} I_H = & \int_a^t \left\{ \frac{d}{dt} [H + (\alpha - 1)p_k \cdot \partial_{n+k} H - \partial_0 H \cdot \right. \\ & \left. \mu(\tau)] - \partial_0 H \right\} \cdot \xi_{H0} d\tau - [H + (\alpha - 1)p_k \cdot \right. \\ & \left. \partial_{n+k} H - \partial_0 H \cdot \mu(t)] \cdot \xi_{H0} + \int_a^t (p_k \cdot \right. \\ & \left. {}^C D_\gamma^{\alpha,\beta} \xi_{Hk} + \xi_{Hk}^\sigma \cdot {}^{RL} D_{\Delta,1-\gamma}^{\beta,\alpha} p_k) d\tau \end{aligned} \quad (71)$$

定理 5 对于联合 Caputo 分数阶 Hamilton 系统 [式(41)], 若无限小生成元满足式(70), 则该系统存在形如式(71)的守恒量.

注 16 当 $\alpha \rightarrow 1, \beta \rightarrow 1$ 时, 对于时间尺度上的 Hamilton 系统[式(42)], 其 Noether 恒等式为

$$p_j \cdot \xi_{Hj}^\Delta - H \xi_{H0}^\Delta - \partial_0 H \xi_{H0} - \partial_k H \cdot \xi_{Hk}^\sigma = 0 \quad (72)$$

对应 Noether 守恒量为

$$\begin{aligned} I_H = & \int_a^t \{ [H - \partial_0 H \cdot \mu(\tau)]^\Delta - \partial_0 H \} \cdot \xi_{H0}^\sigma \Delta\tau - \\ & [H - \partial_0 H \cdot \mu(t)] \cdot \xi_{H0} + p_k \cdot \xi_{Hk} \end{aligned} \quad (73)$$

此结论与文献[45]所得结果一致.

注 17 当 $\gamma = 1$ 时, 对于时间尺度左 Caputo 分数阶 Hamilton 系统[式(43)], 其 Noether 恒等式为

$$\begin{aligned} p_j \cdot {}^C D_{\Delta,t}^\alpha \xi_{Hj} - (\alpha - 1)p_j \cdot \xi_{H0}^\Delta {}^C D_{\Delta,t}^\alpha q_j - \\ H \xi_{H0}^\Delta - \partial_0 H \cdot \xi_{H0} - \partial_k H \cdot \xi_{Hk}^\sigma = 0 \end{aligned} \quad (74)$$

对应 Noether 守恒量为

$$I_H = \int_a^t \{ [H + (\alpha - 1)p_k \cdot \partial_{n+k} H - \partial_0 H \cdot$$

$$\begin{aligned} \mu(\tau)]^\Delta - \partial_0 H \} \cdot \xi_{H0}^\sigma \Delta\tau - [H + (\alpha - 1)p_k \cdot \\ \partial_{n+k} H - \partial_0 H \cdot \mu(t)] \cdot \xi_{H0} + \int_a^t (p_k \cdot {}^C D_{\Delta,t}^\alpha \xi_{Hk} - \\ \xi_{Hk}^\sigma \cdot {}^{RL} D_{\Delta,b}^\alpha p_k) \Delta\tau \end{aligned} \quad (75)$$

此结果与文献[39]所得结果一致.

注 18 当 $\gamma = 0$ 时, 对于时间尺度右 Caputo 分数阶 Hamilton 系统[式(44)], 其 Noether 恒等式为

$$\begin{aligned} -p_j \cdot {}^C D_{\Delta,b}^\alpha \xi_{Hj} + (\alpha - 1)p_j \cdot \xi_{H0}^\Delta {}^C D_{\Delta,b}^\alpha q_j - \\ H \xi_{H0}^\Delta - \partial_0 H \cdot \xi_{H0} - \partial_k H \cdot \xi_{Hk}^\sigma = 0 \end{aligned} \quad (76)$$

对应 Noether 守恒量为

$$\begin{aligned} I_H = & \int_a^t \{ [H + (\alpha - 1)p_k \cdot \partial_{n+k} H - \partial_0 H \cdot \\ & \mu(\tau)]^\Delta - \partial_0 H \} \cdot \xi_{H0}^\sigma \Delta\tau - [H + (\alpha - 1)p_k \cdot \\ & \partial_{n+k} H - \partial_0 H \cdot \mu(t)] \cdot \xi_{H0} - \int_a^t (p_k \cdot {}^C D_{\Delta,b}^\alpha \xi_{Hk} - \\ & \xi_{Hk}^\sigma \cdot {}^{RL} D_{\Delta,t}^\alpha p_k) \Delta\tau \end{aligned} \quad (77)$$

注 19 当 $\alpha = \beta, \gamma = 1/2$ 时, 对于时间尺度 Riesz-Caputo 分数阶 Hamilton 系统[式(45)], 其 Noether 恒等式为

$$\begin{aligned} p_j \cdot {}^C D_{\Delta,b}^\alpha \xi_{Hj} - (\alpha - 1)p_j \cdot \xi_{H0}^\Delta {}^C D_{\Delta,b}^\alpha q_j - \\ H \xi_{H0}^\Delta - \partial_0 H \cdot \xi_{H0} - \partial_k H \cdot \xi_{Hk}^\sigma = 0 \end{aligned} \quad (78)$$

对应 Noether 守恒量为

$$\begin{aligned} I_H = & \int_a^t \{ [H + (\alpha - 1)p_k \cdot \partial_{n+k} H - \partial_0 H \cdot \\ & \mu(\tau)]^\Delta - \partial_0 H \} \cdot \xi_{H0}^\sigma \Delta\tau - [H + (\alpha - 1)p_k \cdot \\ & \partial_{n+k} H - \partial_0 H \cdot \mu(t)] \cdot \xi_{H0} + \int_a^t (p_k \cdot {}^C D_{\Delta,b}^\alpha \xi_{Hk} + \\ & \xi_{Hk}^\sigma \cdot {}^{RL} D_{\Delta,b}^\alpha p_k) \Delta\tau \end{aligned} \quad (79)$$

4 算例

例 1 研究分数阶 Kepler 问题. 时间尺度联合 Caputo 分数阶 Lagrange 函数为

$$L(q_1^\sigma, q_2^\sigma, {}^C D_{\Delta,\gamma}^{\alpha,\beta} q_1, {}^C D_{\Delta,\gamma}^{\alpha,\beta} q_2) = \frac{1}{2} [({}^C D_{\Delta,\gamma}^{\alpha,\beta} q_1)^2 + \\ ({}^C D_{\Delta,\gamma}^{\alpha,\beta} q_2)^2] + [(q_1^\sigma)^2 + (q_2^\sigma)^2]^{-\frac{1}{2}} \quad (80)$$

考虑该系统的 Noether 对称性问题.

根据式(25)可得运动微分方程为

$$\begin{aligned} q_1^\sigma [(q_1^\sigma)^2 + (q_2^\sigma)^2]^{-\frac{3}{2}} + {}^{RL} D_{\Delta,1-\gamma}^{\beta,\alpha} {}^C D_{\Delta,\gamma}^{\alpha,\beta} q_1 = 0, \\ q_2^\sigma [(q_1^\sigma)^2 + (q_2^\sigma)^2]^{-\frac{3}{2}} + {}^{RL} D_{\Delta,1-\gamma}^{\beta,\alpha} {}^C D_{\Delta,\gamma}^{\alpha,\beta} q_2 = 0 \end{aligned} \quad (81)$$

根据式(52)可得 Noether 恒等式

$$-\frac{1}{2} [(q_1^\sigma)^2 + (q_2^\sigma)^2]^{-\frac{3}{2}} \cdot 2q_1^\sigma \cdot \xi_{L1}^\sigma -$$

$$\begin{aligned} & \frac{1}{2}[(q_1^\sigma)^2 + (q_2^\sigma)^2]^{-\frac{3}{2}} \cdot 2q_2^\sigma \cdot \xi_{L2}^\sigma + {}^C D_{\Delta,\gamma}^{\alpha,\beta} q_1 \cdot \\ & ({}^C D_{\Delta,\gamma}^{\alpha,\beta} \xi_{L1} - \alpha {}^C D_{\Delta,\gamma}^{\alpha,\beta} q_1 \cdot \xi_{L0}^\Delta) + {}^C D_{\Delta,\gamma}^{\alpha,\beta} q_2 \cdot \\ & ({}^C D_{\Delta,\gamma}^{\alpha,\beta} \xi_{L2} - \alpha {}^C D_{\Delta,\gamma}^{\alpha,\beta} q_2 \cdot \xi_{L0}^\Delta) + \left\{ \frac{1}{2} [({}^C D_{\Delta,\gamma}^{\alpha,\beta} q_1)^2 + \right. \\ & \left. ({}^C D_{\Delta,\gamma}^{\alpha,\beta} q_2)^2] + [(q_1^\sigma)^2 + (q_2^\sigma)^2]^{-\frac{1}{2}} \right\} \xi_{L0}^\Delta = 0 \end{aligned} \quad (82)$$

由式(82)可得解

$$\xi_{L0} = 0, \xi_{L1} = -q_2, \xi_{L2} = q_1 \quad (83)$$

其对应守恒量为

$$\begin{aligned} I_L = & \int_a^t (-q_2^\sigma \cdot {}^{RL} D_{\Delta,1-\gamma}^{\beta,a} {}^C D_{\Delta,\gamma}^{\alpha,\beta} q_1 + q_1^\sigma \cdot \\ & {}^{RL} D_{\Delta,1-\gamma}^{\beta,a} {}^C D_{\Delta,\gamma}^{\alpha,\beta} q_2) d\tau = \text{const.} \end{aligned} \quad (84)$$

当 $\gamma=1$ 时

$$\begin{aligned} I_L = & \int_a^t (q_2^\sigma \cdot {}_r^{RL} D_{\Delta,b}^{\alpha,a} {}^C D_{\Delta,\tau}^{\alpha,\beta} q_1 - \\ & q_1^\sigma \cdot {}_r^{RL} D_{\Delta,b}^{\alpha,a} {}^C D_{\Delta,\tau}^{\alpha,\beta} q_2) d\tau = \text{const.} \end{aligned} \quad (85)$$

此结果与文献[38]所得结果一致.

当 $\alpha \rightarrow 1, \beta \rightarrow 1$ 时

$$\begin{aligned} I_L = & \int_a^t (-q_2^\sigma \cdot q_1^{\Delta\Delta} + q_1^\sigma \cdot q_2^{\Delta\Delta}) d\tau \\ = & -q_2 q_1^\Delta + q_1 q_2^\Delta = \text{const.} \end{aligned} \quad (86)$$

此结果与文献[15]所得结果一致.

例 2 研究单自由度线性分数阶振动系统的 Noether 对称性, 其中 Lagrange 函数为

$$L = \frac{1}{2} e^{\xi t} [({}^C D_{\Delta,\gamma}^{\alpha,\beta} q)^2 - (q^\sigma)^2] \quad (87)$$

时间尺度取 $T = \{a + mh, m \in \mathbb{N}\}$.

由式(87)可得广义动量和 Hamilton 函数为

$$p = e^{\xi t} \cdot {}^C D_{\Delta,\gamma}^{\alpha,\beta} q, H = \frac{p^2}{2e^{\xi t}} + \frac{1}{2} e^{\xi t} (q^\sigma)^2 \quad (88)$$

根据式(40)可得运动微分方程为

$${}^{RL} D_{\Delta,1-\gamma}^{\beta,a} p = -e^{\xi t} \cdot q^\sigma, {}^C D_{\Delta,\gamma}^{\alpha,\beta} q = \frac{2p}{2e^{\xi t}} = \frac{p}{e^{\xi t}} \quad (89)$$

根据式(68)可得 Noether 恒等式

$$\begin{aligned} p \cdot {}^C D_{\Delta,\gamma}^{\alpha,\beta} \xi_H - (\alpha - 1) p \cdot \xi_{H0}^\Delta \cdot {}^C D_{\Delta,\gamma}^{\alpha,\beta} q - \\ \left[\frac{p^2}{2e^{\xi t}} + \frac{1}{2} e^{\xi t} (q^\sigma)^2 \right] \xi_{H0}^\Delta - \left[\frac{1}{2} e^{\xi t} (q^\sigma)^2 \right] \zeta + \\ \frac{1}{2} p^2 \cdot \frac{-\zeta}{e^{\xi t}} \cdot \xi_{H0} - e^{\xi t} q^\sigma \cdot \xi_H^\sigma = 0 \end{aligned} \quad (90)$$

由式(90)可得解

$$\xi_{H0} = 1, \xi_H = -\frac{\zeta}{2} q \quad (91)$$

其对应守恒量为

$$\begin{aligned} I_H = & \frac{1}{2} \int_a^t \left[\frac{\zeta p^2}{e^{\xi t}} - \zeta (q^\sigma)^2 e^{\xi t} \right] d\tau - \\ & \frac{\zeta q}{2} [\gamma {}_r I_{\Delta,b}^{1-a} p + (1-\gamma) {}_a I_{\Delta,t}^{1-\beta} p] \end{aligned} \quad (92)$$

当 $\gamma=1$ 时

$$\begin{aligned} I_H = & \frac{1}{2} \int_a^t \left[\frac{\zeta p^2}{e^{\xi t}} - \zeta (q^\sigma)^2 e^{\xi t} \right] d\tau - \frac{\zeta q}{2} {}_r I_{\Delta,b}^{1-a} p \\ = & \text{const.} \end{aligned} \quad (93)$$

此结果与文献[39]所得结果一致.

5 结论

文中讨论了时间尺度联合 Caputo 分数阶 Lagrange 系统和时间尺度联合 Caputo 分数阶 Hamilton 系统的 Noether 定理, 并给出若干特例. 有些特例的结果与原有结果一致, 有些特例的结论是新的. 广义分数阶导数是更为广泛的导数, 目前时间尺度广义分数阶导数微积分尚未研究, 后续也可考虑时间尺度广义分数阶 Noether 定理.

参考文献

- [1] NOETHER A E. Invariante variationsprobleme [J]. Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen, Mathematisch-Physikalische Klasse, 1918(2): 235–257
- [2] 梅凤翔. 分析力学[M]. 北京: 北京理工大学出版社, 2013.
- [3] MEI F X. Analytical mechanics [M]. Beijing: Beijing Institute of Technology Press, 2013. (in Chinese)
- [4] MEI F X, WU H B. Dynamics of constrained mechanical systems [M]. Beijing: Beijing Institute of Technology Press, 2009.
- [5] MEI F X, WU H B, ZHANG Y F. Symmetries and conserved quantities of constrained mechanical systems [J]. International Journal of Dynamics and Control, 2014, 2(3): 285–303.
- [6] 赵跃宇, 梅凤翔. 力学系统的对称性与不变量[M]. 北京: 科学出版社, 1999.
- ZHAO Y Y, MEI F X. Symmetry and invariants of dynamical systems [M]. Beijing: Science Press, 1999. (in Chinese)
- [7] 梅凤翔. 约束力学系统的对称性与守恒量[M]. 北京: 北京理工大学出版社, 2004.

- MEI F X. Symmetry and conserved quantity of constrained mechanical systems [M]. Beijing: Beijing Institute of Technology Press, 2004. (in Chinese)
- [7] 梅凤翔. 李群和李代数对约束力学系统的应用 [M]. 北京: 科学出版社, 1999.
- MEI F X. Applications of Lie groups and Lie algebras to constrained mechanical systems [M]. Beijing: Science Press, 1999. (in Chinese)
- [8] 董欣畅, 张毅. 非保守非完整系统的 Herglotz 型 Noether 定理[J]. 动力学与控制学报, 2024, 22(5): 1–7.
- DONG X C, ZHANG Y. Herglotz-type Noether theorem for nonconservative nonholonomic system [J]. Journal of Dynamics and Control, 2024, 22(5): 1–7. (in Chinese)
- [9] 黄丽琴, 张毅. 二阶非完整系统 Vacco 动力学的 Herglotz 型 Noether 定理[J]. 动力学与控制学报, 2024, 22(9): 16–23.
- HUANG L Q, ZHANG Y. Herglotz-type Noether theorem for Vacco dynamics of second-order non-holonomic systems [J]. Journal of Dynamics and Control, 2024, 22(9): 16–23. (in Chinese)
- [10] HILGER S. Ein maßkettenkalkül mit anwendung auf zentrumsmannigfaltigkeiten [D]. Würzburg, Germany: Universität Würzburg, 1988.
- [11] BOHNER M, PETERSON A C. Dynamic equations on time scales: an introduction with applications [M]. Boston: Birkhäuser, 2001
- [12] BOHNER M, PETERSON A. Advances in dynamic equations on time scales [M]. Boston: Birkhäuser, 2003.
- [13] BARTOSIEWICZ Z, TORRES D F M. Noether's theorem on time scales [J]. Journal of Mathematical Analysis and Applications, 2008, 342(2): 1220–1226.
- [14] MARTINS N, TORRES D F M. Noether's symmetry theorem for nabla problems of the calculus of variations [J]. Applied Mathematics Letters, 2010, 23(12): 1432–1438.
- [15] ANEROT B, CRESSON J, BELGACEM K H, et al. Noether's s-type theorems on time scales [J]. Journal of Mathematical Physics, 2020, 61(11): 113502.
- [16] ZHANG Y. Nonshifted dynamics of constrained systems on time scales under Lagrange framework and its Noether's theorem [J]. Communications in Nonlinear Science and Numerical Simulation, 2022, 108: 106214.
- [17] PENG K K, LUO Y P. Dynamics symmetries of Hamiltonian system on time scales [J]. Journal of Mathematical Physics, 2014, 55(4): 042702.
- [18] 张毅. 时间尺度上 Hamilton 系统的 Noether 理论 [J]. 力学季刊, 2016, 37(2): 214–224.
- ZHANG Y. Noether theory for Hamiltonian system on time scales [J]. Chinese Quarterly of Mechanics, 2016, 37(2): 214–224. (in Chinese)
- [19] SONG C J, ZHANG Y. Conserved quantities for Hamiltonian systems on time scales [J]. Applied Mathematics and Computation, 2017, 313: 24–36.
- [20] SONG C J, ZHANG Y. Noether theorem for Birkhoffian systems on time scales [J]. Journal of Mathematical Physics, 2015, 56(10): 102701.
- [21] SONG C J, ZHANG Y. Noether theory for Birkhoffian systems with nabla derivatives [J]. Journal of Nonlinear Sciences and Applications, 2017, 10(4): 2268–2282.
- [22] HOU S, SONG C J. Perturbation to symmetries and adiabatic invariants for generalized Birkhoffian systems on time scales [J]. Wuhan University Journal of Natural Sciences, 2024, 29(3): 263–272.
- [23] CAI P P, FU J L, GUO Y X. Noether symmetries of the nonconservative and nonholonomic systems on time scales [J]. Science China: Physics, Mechanics & Astronomy, 2013, 56(5): 1017–1028.
- [24] ZU Q H, ZHU J Q. Noether theorem for nonholonomic nonconservative mechanical systems in phase space on time scales [J]. Journal of Mathematical Physics, 2016, 57(8): 082701.
- [25] LUO S K, XU Y L. Fractional Birkhoffian mechanics [J]. Acta Mechanica, 2015, 226(3): 829–844.
- [26] FREDERICO G S F, TORRES D F M. A formulation of Noether's theorem for fractional problems of the calculus of variations [J]. Journal of Mathematical Analysis & Applications, 2007, 334(2): 834–846.
- [27] ZHOU S, FU H, FU J L. Symmetry theories of Hamiltonian systems with fractional derivatives [J]. Science China: Physics, Mechanics & Astronomy, 2011, 54(10): 1847.
- [28] JIN S X, ZHANG Y. Noether symmetries for non-conservative Lagrange systems with time delay based on fractional model [J]. Nonlinear Dynamics, 2015, 79(2): 1169–1183.
- [29] ZHAI X H, ZHANG Y. Noether symmetries and

- conserved quantities for fractional Birkhoffian systems with time delay [J]. Communications in Nonlinear Science & Numerical Simulation, 2016, 36: 81—97.
- [30] FREDERICO G S F, TORRES D F M. Fractional optimal control in the sense of Caputo and the fractional Noether's theorem [J]. International Mathematical Forum, 2008, 3(10): 479—493.
- [31] FREDERICO G S F, TORRES D F M. Fractional Noether's theorem in the Riesz-Caputo sense [J]. Applied Mathematics & Computation, 2010, 217 (3): 1023—1033.
- [32] SONG C J, ZHANG Y. Noether symmetry and conserved quantity for FractiOnal birkhoffian mechanics and its applications [J]. Fractional Calculus & Applied Analysis, 2018, 21(2): 509—526.
- [33] SONG C J, ZHAI X H. Noether theorem for fractional singular systems [J]. Wuhan University Journal of Natural Sciences, 2023, 28(3): 207—216.
- [34] ANASTASSIOU G A. Principles of delta fractional calculus on time scales and inequalities [J]. Mathematical and Computer Modelling, 2010, 52(3/4): 556—566.
- [35] BENKHETTOU N, HAMMOUDI A, TORRES D F M. Existence and uniqueness of solution for a fractional Riemann-Liouville initial value problem on time scales [J]. Journal of King Saud University-Science, 2016, 28(1): 87—92.
- [36] WU G C, BALEANU D. Discrete fractional logistic map and its chaos [J]. Nonlinear Dynamics, 2014, 75(1): 283—287.
- [37] HUANG L L, PARK J H, WU G C, et al. Variable-order fractional discrete-time recurrent neural networks [J]. Journal of Computational and Applied Mathematics, 2020, 370: 112633.
- [38] 田雪, 张毅. Caputo Δ 型分数阶时间尺度 Noether 定理[J]. 力学学报, 2021, 53(7): 2010—2022.
- TIAN X, ZHANG Y. Caputo Δ -type fractional time-scales Noether theorem [J]. Chinese Journal of Theoretical and Applied Mechanics, 2021, 53(7): 2010—2022. (in Chinese)
- [39] TIAN X, ZHANG Y. Fractional time-scales Noether theorem with Caputo Δ derivatives for Hamiltonian systems [J]. Applied Mathematics and Computation, 2021, 393: 125753.
- [40] TIAN X, ZHANG Y. Caputo Δ -type fractional time-scales Noether theorem of Birkhoffian systems [J]. Acta Mechanica, 2022, 233(11): 4487—4503.
- [41] GEORGIEV S G. Fractional dynamic calculus and fractional dynamic equations on time scales [M]. Cham Switzerland: Springer, 2018.
- [42] BAHAA G M, TORRES D F M. Time-fractional optimal control of initial value problems on time scales [M]//AREA I, CABADA A, CID J A, et al. Nonlinear Analysis and Boundary Value Problems. Cham: Springer, 2019: 229—242.
- [43] ZHANG Y. Fractional differential equations of motion in terms of combined Riemann-Liouville derivatives [J]. Chinese Physics B, 2012, 21 (8): 084502.
- [44] 张毅. 时间尺度上完整非保守力学系统的 Noether 定理[J]. 苏州科技大学学报(自然科学版), 2020, 37(1): 6—11.
- ZHANG Y. Noether's theorem for holonomic non-conservative mechanical systems on time scales [J]. Journal of Suzhou University of Science and Technology (Natural Science), 2020, 37(1): 6—11. (in Chinese)
- [45] 宋传静, 侯爽. 时间尺度上约束力学系统的 Noether 型绝热不变量[J]. 力学学报, 2024, 56 (8): 2397—2407.
- SONG C J, HOU S. Noether-type adiabatic invariants for constrained mechanics systems on time scales [J]. Chinese Journal of Theoretical and Applied Mechanics, 2024, 56 (8): 2397—2407. (in Chinese)